

Supersymmetry and Identical Bands

P. von Brentano

Institut für Kernphysik, Universität zu Köln, Germany

Supersymmetry as applied to identical bands is discussed. A review of the work of the Koeln-Dubna group on this topic is given and examples in $^{171,172}\text{Yb}$, $^{173,174}\text{Hf}$ and $^{195,194}\text{Pt}$ are discussed. The role of pseudo-spin in the supersymmetry is investigated. A recent precision lifetime measurement for identical bands in $^{171,172}\text{Yb}$ is discussed. Keywords: U(6/12) supersymmetry, identical bands, pseudo-spin

I. INTRODUCTION

Supersymmetry has been successfully transferred from particle physics to nuclear physics by Franco Iachello and coworkers[1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11]. Various attempts to obtain an extended classification, in which several neighboring nuclei including even, odd and odd-odd ones are described by the same Hamiltonian, have led to convincing results e.g.[12]. However, in nuclear structure theory primary attention has been paid to the investigation of models with Hamiltonians which exhibit dynamical supersymmetries. In a recent paper[13] it has been shown, however, that the concepts of identical bands[14, 15, 16] and pseudo-spin [18, 19] allows one to discuss (partial) supersymmetry also for cases in which the even partner nucleus has no dynamical symmetry. This has also been discussed in the consistent Q formalism[21, 22]. The concept of “Identical Bands” (IB) was suggested by a Strasbourg-Liverpool collaboration led by Byrski [14], who found an agreement to better than 0.3% in the gamma energies of two superdeformed

(sd) bands in ^{151}Tb and ^{152}Dy . The role of pseudo-spin in identical bands was subsequently clarified by Nazarewicz et al. [15]. An explanation based on the $U(6/12)$ supersymmetry was given in the frame of the Interacting Boson Model by Gelberg et al. [23] and in the frame of supersymmetric quantum mechanics by Amado, Bijker et al. [24]. Although the importance of the discovery of the IB was widely recognized the data on the Dy-Tb pair are still incomplete due to experimental difficulties. The group at Argonne has recently settled the spins of the identical superdeformed band in ^{152}Dy [17] but the spins in ^{151}Tb are still unknown. Also the $3/2, 7/2\dots$ signature partner band in ^{151}Tb is still missing. This shows that one has to rely on the spectroscopy of “Identical Bands” at normal deformation, where spins and parities are all known. In this case the moments of inertia of the even and odd nucleus system tend to differ by 10% or more as compared to 0.3% in the SD case. A survey is given by Baktash et al.[16]. We show in Fig. 1 a convincing example of the identical bands in ^{174}Hf and ^{173}Hf [13].

In the following I will review the Koeln-Dubna work[13, 25, 26, 27] which focused on the supersymmetry connected with identical bands. In order to avoid equations or technical discussions a simple schematic model for supersymmetry will be discussed. This is done on the basis of the $U(6/12)$ supersymmetry [1, 7] which has a very transparent microscopic foundation. In $U(6/12)$ respectively in $U(15/30)$ [1, 7] there are three (five) contributing shells in the odd nucleus and which have the spins $1/2, 3/2$ and $5/2$ (or $1/2, 3/2, 5/2, 7/2$ and $9/2$)

II. PSEUDO-SPIN

In discussing the $U(6/12)$ SUSY and the schematic model pseudo-spin plays an important role. Thus I make a few comments on pseudo-spin symmetry. This approximate symmetry is long known [18, 19]. Recently it was shown, that pseudo-spin symmetry is actually based on a relativistic symmetry[20]. We remind briefly




Fig.1

Comparison of the energies of SUSY partner bands in ^{174}Hf and ^{173}Hf . The energies labeled CALC are obtained by correcting for the change in the moment of inertia and by allowing a pseudo-spin orbit coupling. From Jolos et al.[13].

the pseudo-spin scheme: In Fig. 2 the s.p. neutron states around ^{208}Pb are shown. In the pseudo-spin scheme the $3p_{1/2}$, $2f_{5/2}$ and $3p_{3/2}$ levels are relabeled with the pseudo-spin quantum numbers $2\tilde{s}_{1/2}$, $2\tilde{d}_{5/2}$, $2\tilde{d}_{3/2}$. Whereas the angular momenta of the particle are $l = 1$ or $l = 3$ respectively, the pseudo angular momenta are $\tilde{l} = 0$ or $\tilde{l} = 2$. For simplicity it is assumed in the following that the pseudo-spin of the last odd particle is fully decoupled in energy. In this case to each state with spin I_e of the even nucleus there belongs a multiplet of states in the odd partner nucleus with the same excitation energy and with spins :




Fig. 2

The figure shows on the right side the spins of the neutron single particle energies in Pb and on the left side their pseudo angular momentum: $\tilde{l}$.

$$I_{odd}^+ = I_e + 1/2 \ \& \ I_{odd}^- = | I_e - 1/2 |$$

respectively. In particular there is a doublet of states for $I_e \neq 0$ and a singlet for $I_e = 0$. An example of nearly decoupled pseudo-spin multiplets is found in very nice data by Jolie and Graw[32]. They measured a set of states which is presumably complete below 1 MeV. After this they were able to assign the observed levels to pseudo-spin multiplets. Indeed - as shown in Fig. 3 - each level in ^{195}Pt (below 0.8 MeV) comes either as a doublet or as a singlet if $I_e = 1/2$. This is to -my knowledge - the only case in which pseudo-spin multiplets in the odd nucleus are seen so clearly. So more data are needed. It is a beautiful prediction from the dynamical $U(6/12)$ supersymmetry and from the pseudo-spin concept.

III. SUPERSYMMETRY WITH IDENTICAL BANDS : A SCHEMATIC MODEL.

I will now discuss a schematic model for supersymmetry (SUSY) with identical bands in the frame of $U(6/12)$. The odd particle is in shells with spins $1/2$, $3/2$ and $5/2$ only. The pseudo angular momenta $\tilde{l}$ of the odd particle have integer values and can be considered as bosonic excitations. They have the values $\tilde{l} = 0$ and $\tilde{l} = 2$ corresponding to the spins $L = 0$ and $L = 2$ of the s and d bosons. Thus it is “natural” following T. Otsuka [30] to describe the particles with pseudo angular momenta $\tilde{l} = 0$ and $\tilde{l} = 2$ by (pseudo angular momentum) quasi bosons with the spins $\tilde{l}=0 : s^f$ and $\tilde{l}=2 : d^f$. Thus there are three kinds of bosons : neutron bosons b^n , proton bosons b^p , and quasi bosons b^f

The schematic model assumes :

- Equal energies $\varepsilon_d^n = \varepsilon_d^p = \varepsilon_d^f$ of the three kinds of bosons : b^n , b^p , b^f
- Equal interactions V_{ik} among the three kinds of bosons b^n , b^p , b^f .
- Fully decoupled pseudo-spin.

These assumptions imply identical bands in the following three bosonic systems : K_0 , K_1 and K_2 defined in Fig. 3 and which have the same total boson numbers N_B : but with different numbers of bosons of the three kinds : b^n , b^p , b^f .

In particular one finds :

- to every state in K_0 there corresponds a state in K_1 with the same energy.
 K_0 and K_1 are members of an F spin multiplet. They have good F-spin.
- to every state in K_1 there corresponds a state in K_2 with the same energy
 and vice versa
- to every state in K_2 there corresponds a pseudo-spin multiplet in K_3 with the same energy

Identical Bands in 4 Systems




Fig. 3

The figure illustrates the “Identical Bands” suggested by the schematic model in four “nuclei” which have the same total number of “bosons” but differing numbers of bosons of various types as shown in the figure. The “nucleus” K_3 is obtained from K_2 by adding the pseudo-spin $\tilde{s}$

In summary in the schematic model to each pseudo-spin multiplet of the odd system there belongs a unique state of the corresponding even system with the same energy. This is a somewhat surprising result. It means that there are as many states in the even “nucleus” K_1 as there are pseudo-spin multiplets in the “odd nucleus” K_3 . One might think that the odd nucleus has many more states. One must remember, however, that also the “mixed symmetry” states in the “even nucleus” have to be considered.

IV. COMPARISON OF SCHEMATIC MODEL TO DATA

Below the following predictions A)... D) of the schematic model are compared to data:

- A) to each level below about 0.8 MeV in ^{194}Pt there is a multiplet in ^{195}Pt with similar energy. We note that not only the yrast band but e.g. also the gamma band has a SUSY partner. (see Fig. 4 [32]) We note that the energies of the centroids of the doublets differ somewhat from the energy of the corresponding level in ^{194}Pt . Also there is some splitting in the doublets. Thus the schematic model is broken somewhat
- B) There are low lying multiplets in ^{195}Pt which do not correspond to low lying states in ^{194}Pt , however. In the schematic model they might correspond to high lying mixed symmetry states [31]. An example is the (199 keV and 224 keV) doublet, which has a core state with $I^\pi = 1^+$.
- C) in the rare earth region there are identical bands with a vanishing projection $\widetilde{L}_3 = \widetilde{\Lambda} = 0$ of the pseudo angular momentum on the symmetry axis. The $\widetilde{\Lambda} = 0$ band qualifies as super symmetrical partner band of the $K = 0$ ground band of the even partner nucleus. These were the bands considered in [13, 26, 27]. Examples are identical bands in $^{171,172}\text{Yb}$ (Fig. 5), $^{173,174}\text{Hf}$ (Fig. 1) and in $^{181,2}\text{Pt}$ [26]. These are convincing SUSY partner bands.
- D) the bands in $^{171,172}\text{Yb}$, in $^{173,174}\text{Hf}$ and in $^{181,1822}\text{Pt}$ which have a non vanishing pseudo angular projection $\widetilde{L}_3 = \widetilde{\Lambda} \neq 0$ have a pseudo-spin partner band which is separated in energy by a few hundred keV or more. Thus pseudo-spin is not decoupled for these bands. And the schematic model breaks down for these bands.

We note that A) and C) show that there are identical bands and spectra. B) indicates that there are levels in these nuclei which have no partners with the same energies. Thus the schematic SUSY survives only for a part of the levels as a partial symmetry[28, 29]. We note, however, that these extra states are reproduced in the dynamical U(6/12) SUSY as this uses a different (more correct) interaction. Part of the reason for the failure is certainly that pseudo-spin is not fully decoupled i.e. the excitation energies in the pseudo-spin doublets are not equal. In ^{207}Pb one finds : $\Delta E (2f5/2 - 3p3/2) = 329\text{keV}$. However the pseudo-spin splitting is still much smaller than the spin orbit splitting : $\Delta E (2f5/2 - 3p3/2) \leq 0.36 * \Delta E (3p1/2 - 3p3/2)$. An interesting point concerns the apparent validity of pseudo-spin in the $\tilde{\Lambda} = 0$ bands in $^{171,172}\text{Yb}$, $^{173,174}\text{Hf}$ and $^{181,182}\text{Pt}$ (see C) and the fact that pseudo-spin is not decoupled for the $\tilde{\Lambda} \neq 0$. The reason may be that the $\tilde{\Lambda} = 0$ state has no pseudo-spin partner, as is shown e.g. in Fig. 2 for the $3p1/2$ state, whereas for $\tilde{\Lambda} \neq 0$ there are two pseudo-spin partner levels e.g. the $2f5/2$ and $3p3/2$ levels, which can interact. The small decoupling in the $\tilde{\Lambda} = 0$ bands is also interesting and deserves a closer study.

V. SUPERSYMMETRY OF IDENTICAL BANDS IN $^{171,172}\text{Yb}$ SUPPORTED BY LIFETIME DATA

There are 5 requirements for an $K = 0$ and an odd rotational band in neighbor nuclei to be super partners of an unbroken partial supersymmetry:

- 1) identity of the moments of inertia,
- 2a) decoupling of the pseudo-spin.(close doublets in the odd nucleus),
- 2b) alternatively a special value $a=1$ of the decoupling parameter in the rotational energy formula of the $K = 1/2$ band,
- 3) an even pseudo angular momentum of the particle,
- 4) identity of the quadrupole transition moments, and

5) $K = 1/2$, $\tilde{\Lambda}=0$ in the odd band.

Some people like to remember these 5 conditions and these special values of special parameters. Others like to say there is a SUSY, that implies all.

At this point it might be appropriate to avoid a possible misunderstanding, namely that the $K = 1/2$, $\tilde{\Lambda}=0$ bands e.g. in ^{171}Yb could be obtained by a weak coupling of the spin of the core state I of ^{172}Yb to the pseudo-spin.

$$J = I \oplus \tilde{s}$$

This is not true at all. Rather one couples the spins I of several core states strongly to the pseudo angular momentum $\tilde{l}$ of the particle and after this the pseudo-spin is weakly coupled.

$$J = \Sigma(I, \tilde{l}) [(I \oplus \tilde{l}) \oplus \tilde{s}]$$

Thus neither I nor $\tilde{l}$ are not even approximately good quantum numbers. And although the $^{171,172}\text{Yb}$ pair looks like a weak coupling situation, it is actually a quite complicated coupling scheme: strong for $\tilde{l}$ and weak for $\tilde{s}$ [26].

Thus it is important to test the SUSY in more detail by establishing the identity of the quadrupole transitions. The pair $^{171,172}\text{Yb}$ is the only case in which one do Coulomb Excitation on both for both SUSY partners. The energies are shown in Fig. 5 agree with the SUSY. For these reasons Klug, Dewald et al. measured precision lifetimes by the recoil distance method following Coulomb excitation experiments using an EUROBALL CLUSTER detector at the FN-Tandem accelerator in Koeln [27]. From these data and previous data on branching and mixing ratios from Canberra, the ratios of the transition quadrupole moments for corresponding E2 transitions in ^{171}Yb and ^{172}Yb were obtained , which are given in the Table 1.

One notes that the values of the transition quadrupole moment Q_t for ^{171}Yb and ^{172}Yb are in equal within an error of a few percent[27]. This is one of the most accurate experimental comparisons of transition quadrupole moments for

Tab.1

Ratios of transition quadrupole moments Q_t from SUSY partner bands in ^{171}Yb and ^{172}Yb . The Q_t values are obtained from the measured $B(E2)$ values by using the formulas for the quadrupole moments for the rigid rotor. From Klug, Dewald et al.[27]

	$\Delta I=2$	$\Delta I = 2$	$\Delta I = 1$
I	$\frac{Q_t^{171}(I+1/2)}{Q_t^{172}(I)}$	$\frac{Q_t^{171}(I-1/2)}{Q_t^{172}(I)}$	$\frac{Q_t^{171}(I-1/2)}{Q_t^{172}(I)}$
8	1.02(6)	1.03(6)	1.21(10)
6	0.94(5)	0.94(5)	0.95(6)
4	0.96(2)	0.95(3)	0.97(2)

supersymmetry partner bands. This striking agreement supports strongly the proposed supersymmetry for ^{171}Yb and ^{172}Yb . Indeed all 5 of the requirements for a supersymmetry are fulfilled for the ^{171}Yb and ^{172}Yb pair. On the contrary in the ^{171}Tm and ^{172}Yb pair only three conditions are fulfilled, whereas the conditions 3) and 5): an even pseudo angular momentum of the particle are violated. And indeed there is no supersymmetry in this case.

Summing up convincing examples of (partial) SUSY for energies and $B(E2)$ values in identical bands are shown. An intuitive understanding of the $U(6/12)$ SUSY in the frame of the pseudo-spin concept and the schematic model have been given. Limitations of supersymmetry and pseudo-spin symmetry restricting the SUSY to a partial SUSY have been discussed.


Acknowledgments

My particular thanks go to Franco Iachello, for his constant inspiration to Koeln work since many years. Thanks go also to A. Dewald, A. Gelberg, J. Jolie, R. V. Jolos and T. Otsuka for joint works published and in progress on supersymmetry which were used in this talk. Thanks for many discussions to R. F. Casten, F. Iachello, P. van Isacker and N. Pietralla. This work was partially supported by

the DFG under the contract Br 799/10-2.

-
- [1] F. Iachello and P. Van Isacker, *The Interacting Boson–Fermion Model*, (Cambridge University Press, Cambridge, 1991).
 - [2] A. Frank and P. Van Isacker, *Algebraic Methods in Molecular and Nuclear Physics*, (John Wiley & Sons, New York, 1994).
 - [3] F. Iachello, Phys. Rev. Lett. **44**, 772 (1980).
 - [4] F. Iachello and S. Kuyucak, Ann. Phys. (N.Y.) **136**, 19 (1981).
 - [5] A. B. Balantekin, I. Bars and F. Iachello, Nucl. Phys. **A370**, 284 (1981).
 - [6] J. Jolie, P. Van Isacker, K. Heyde, and A. Frank, Phys. Rev. Lett. **55**, 653 (1985).
 - [7] P. Van Isacker, A. Frank and Hong-Zhou Sun, Ann. Phys.(N.Y.) **157** (1984)183
 - [8] R. Bijker and F. Iachello, Ann. Phys. (N.Y.) **161**, 360 (1985).
 - [9] D. D. Warner, R. F. Casten, and A. Frank, Phys. Lett. B, **180**, 207 (1986).
 - [10] J. Vervier, P. Van Isacker, J. Jolie, V. K. B. Kota, and R. Bijker, Phys. Rev. C **32**, 1406 (1985).
 - [11] P. Van Isacker, J. Jolie, K. Heyde, and A. Frank, Phys. Rev. Lett. **54**, 653 (1985).
 - [12] A. Metz, J. Jolie, G. Graw, R. Hertenberger, J. Gröger, C. Günter, N. Warr, and Y. Eisermann, Phys. Rev. Lett. **83**, 1542 (1999).
 - [13] R. V. Jolos and P. von Brentano, Phys. Rev. C **60**, 064318 (1999).
 - [14] T. Byrski, F. A. Beck, D. Curien, C. Schuck et al., Phys.Rev.Lett. **64**, 1650 (1990).
 - [15] W. Nazarewicz, P. J. Twin, P. Fallon, and J. D. Garrett, Phys.Rev.Lett., **64**, 1654 (1990).
 - [16] C. Baktash, B. Haas, W. Nazarewicz, Annu.Rev.Nucl.Part.Sci., **45**, 485 (1995)R. Janssens, private communication
 - [17] R. Janssens, private communication.
 - [18] A.Arima, M.Harvey, and K.Shimizu, Phys.Lett.,**30 B**, 517 (1969).

- [19] K.T.Hecht and A.Adler Nucl.Physics A 137 129 (1969).
- [20] Ginocchio J N Phys. Rev. Lett. **78**,436 (1997)
- [21] D. D. Warner, R. F. Casten Phys. Rev. C **28**,1798, (1983).
- [22] A. Frank, P. Van Isacker, D. D. Warner, Phys. Lett., **179B**, 474,(1987).
- [23] A. Gelberg, P. von Brentano, R. F. Casten, J.Phys. G **16**, L143 (1990).
- [24] R. D. Amado, R. Bijker, F. Cannata, and J. P. Dedonder, Phys. Rev. Lett., **67**, 2777 (1991).
- [25] R.V.Jolos, P. von Brentano, A. Gelberg, K.-H. Kim, and T.Otsuka, Phys. Lett. B, **430**, 1 (1998).
- [26] R. V. Jolos and P. von Brentano, Phys. Rev. C **63**02, 024304 (2001).
- [27] T. Klug, A. Dewald, R.V. Jolos , B. Saha, P. von Brentano and J.Jolie. Phys. Lett. B 525, 252 (2002).
- [28] Y. Alhassid and A. Leviatan, J. Phys. A **25**, L1265 (1992).
- [29] A. Leviatan, Phys. Rev. Lett. **77**, 818 (1996).
- [30] T.Otsuka, private communication
- [31] P. von Brentano and A. Gelberg, to be published.
- [32] A. Metz, Y. Eisermann, A. Gollwitzer, R. Hertenberger, G. Graw and J. Jolie, Phys. Rev. C 61 064313 (2000).






Fig. 5

The figure compares $K = 1/2$ bands in ^{171}Yb and ^{171}Tm with the $K = 0$ ground band in

^{172}Yb . The bands in Yb have $\tilde{\Lambda} = 0$ and thus form a SUSY doublet. The band in ^{171}Tm has $\tilde{\Lambda} = 1$ (doublet for groundstate) and is thus not the SUSY partner of ^{172}Yb ,

which has $\tilde{\Lambda} = 0$. From Klug et al.[27]